

Prostate Cancer: Serum and Tissue Markers

Gary J. Miller, MD, PhD,* Michael K. Brawer, MD,† Wael A. Sakr, MD,† J. Brantley Thrasher, MD,§ Ronald Townsend, MD*

*University of Colorado Health Sciences Center, Denver, CO; †Northwest Prostate Institute, Seattle, WA;

‡Wayne State University School of Medicine, Detroit, MI; §University of Kansas Medical Center, Kansas City, KS

The detection of prostate cancer, its clinical staging, and the prediction of its prognosis remain topics of paramount importance in clinical management. The digital rectal exam, although once the “gold standard,” has been largely supplanted by a variety of techniques including serum and tissue-based assays. This article reviews recent progress in the development of prostate-specific antigen assays with greater specificity; molecular markers for prostate cancer (DNA ploidy, nuclear morphometry, markers of proliferation, and cell adhesion molecules); the link between vitamin D deficiency and the clinical emergence of prostate cancer; the possible correlation of serum insulin-like growth factor levels with the risk for developing prostate cancer; and the latest advances in radiologic staging. [Rev Urol. 2001;3(suppl 2):S11–S19]

© 2001 MedReviews, LLC

Key words: Prostate-specific antigen • DNA ploidy • E-cadherin • Vitamin D receptors • Insulin-like growth factor • Positron emission tomography • Radiologic staging

The detection of prostate cancer, its clinical staging, and the prediction of its prognosis remain topics of paramount importance in clinical management. The digital rectal exam, although once the “gold standard,” has been largely supplanted by a variety of techniques including serum and tissue-based assays. Methods for analysis of PSA have been improved, and a host of other tumor markers and biologic determinants are on the horizon. Advances in body imaging have also provided new capability for noninvasive assessment.

PSA: An Update

Nothing has been more helpful to advance the management of men with prostatic carcinoma than the development of reliable assays for prostate specific antigen (PSA). Although this analyte was first exploited as a tool to identify prostate cancer in tissue sections, the

suffer from sampling bias, and both PSA velocity and the free/total ratio suffer from test variability.

The recognition that PSA circulates in a number of molecular forms has provided significant enhancement in specificity. Most of PSA in serum is not in the free form found in the ejaculate, but rather it is complexed

atic.^{11,12} These issues make the free/total PSA ratio less useful in broad-based clinical settings.

Because of the concerns over the practicality of the free/total PSA assay, much effort has been expended in attempting to develop specific assays for the complexed form of PSA. This approach is derived from the fact that the ACT complexed form of PSA is the analyte that is more specific for prostatic carcinoma. The Bayer Corporation developed a specific assay for complex PSA that has been shown to be highly reproducible as well as specific.¹³ A study was performed on archival serum utilizing this assay in men undergoing ultrasound guided needle biopsy.¹⁴ In comparison with the Hybritech Tandem R free and total assays, enhanced specificity was found with the Bayer complex form of PSA. For example, at the 95% sensitivity level the specificity of the total PSA was 22%, that of the free/total PSA ratio 15.6% (at a 28% cutoff), and the complexed PSA ratio with a cutoff of 2.52

The greatest liability with PSA, according to most experts, is its relative lack of specificity.

application with the greatest impact has been in early detection and/or screening, where it is used to evaluate risk potential for the presence of malignancy. Despite the tremendous advantages of this, the most important of all human tumor markers, PSA is not an ideal test. Problems, including lack of sensitivity and specificity and the risk of over-detection, are real. The greatest liability with PSA, according to most experts, is its relative lack of specificity. Thus, tremendous efforts have been made to make PSA more specific. Lack of specificity and the inherent false-positive rate is expensive in an economic sense, because it mandates further testing, including antibiotics, ultrasound biopsy, and pathology charges. Moreover, it is expensive emotionally: telling a man that he has an elevated PSA causes a great deal of worry for him as well as for his loved ones.

In an effort to make PSA more specific, so-called PSA derivatives, including PSA velocity, age-specific PSA, and PSA density have been realized. Although there has been considerable enthusiasm for these approaches, none have been shown in broad-based clinical trials to be useful in early detection. Age-specific cutoffs suffer from a lack of sensitivity, PSA density and transition zone density

to protease inhibitors. These include alpha-2 macroglobulin and alpha-1 antichymotrypsin (ACT).¹⁻⁴ The majority of PSA in the circulation is complexed to ACT. Where the complex is formed has not yet been determined. Scandinavian investigators demonstrated that measurement of the free/total ratio of PSA increased the specificity of total PSA alone.^{2,5} Subsequent studies^{6,7} confirmed these findings. The definitive study in this regard was that reported by Catalona et al.⁸ For example, in this multicenter

The potential benefits of utilizing complex PSA determination include cost efficiency when compared with the free/total PSA measurement, which requires two separate PSA determinations.

trial, at the 95% sensitivity level, using a cutoff of 25% led to an enhancement of test specificity of 20% over that afforded by total PSA.

Several problems exist, however, when trying to determine the free/total PSA ratio. There are significant concerns over pre-analytic handling of the specimens, because free PSA is less stable than complexed PSA.^{9,10} Moreover, different manufacturers' assays have given staggeringly different results. The quotient necessary for this ratio makes the bias particularly problem-

afforded a 26.7% specificity. Similar results were obtained from an expanded multi-institutional study.¹⁵ The potential benefits of utilizing complex PSA determination include the use of a single analyte on an elevated platform, which obviates assay variability issues, as well as cost efficiency when compared with the free/total PSA measurement, which requires two separate PSA determinations. Because the complexed PSA assay is equivalent to total PSA for monitoring or staging disease, it would thus appear to be the best

possible form of PSA assay for all clinical uses.

Molecular Markers for Prostate Cancer

In addition to the well-tested clinical utility of stage, grade, and PSA in assessing prognosis and guiding management, the available array of

mous and cannot be exhaustively reviewed in this summary. For the sake of brevity, only DNA ploidy, nuclear morphometry, markers of proliferation, and cell adhesion molecules will be considered.

The DNA content of tumor cells is referred to as *DNA ploidy*. Assessment of DNA ploidy using

nuclear morphometry) have also been reported to be of prognostic significance. In a study of T stage, Gleason score, and 10 nuclear morphometric factors (such as mean nuclear area, nuclear perimeter, shortest and longest nuclear axis, form factor, and their standard deviations), nuclear area had independent prognostic significance, but only in T1 tumors.²⁵ Studies on the technical aspects of these methods have suggested that the superior fixation that occurs in needle biopsy specimens may provide increased accuracy of measurement.²⁶ However, the widespread application of these techniques is plagued by problems with tissue handling, user dependency, standardization of automated techniques using computer equipment, quality assurance, and quality control.²⁷ With these limitations in mind, however, there is continuing interest in these techniques, especially with respect to surrogate endpoints for chemoprevention studies.²⁸

Studies aimed at assessing the rate of proliferation in prostate carcinomas have used a variety of technical

With respect to pT3 tumors, DNA ploidy has prognostic value in predicting response to hormonal therapy.

biologic markers of prostate cancer continues to expand. Some of these serve only as "tumor markers" that can be used to indicate the presence of the disease. Others are more properly designated "biologic determinants," because they are components of the cell that actually have a role in establishing or maintaining the malignant phenotype. The role of a marker is not always clear with respect to establishing prognosis. Confusion arises when researchers attempt to extrapolate between the clinical and laboratory settings. It is also important to recognize that although we attempt to use molecular markers to assist in evaluating early-stage disease, much of the available information about these molecules comes from observations made on material from patients with advanced-stage disease, because of its more ready accessibility. Modern techniques, such as laser-capture microscopy, may overcome this deficit. In addition, the application of any marker to the clinical setting is an extremely complex issue. The reader is referred, therefore, to the criteria for evaluation of prognostic markers and reviews addressing statistical literature that have previously been published.^{16,17} The array of candidates available to enhance routine measurements of grade and stage is enor-

flow cytometry and/or static imaging has been extensively investigated in prostate cancer.^{18,19} Many studies have indicated that abnormal ploidy (ie, aneuploidy or tetraploidy) correlates with tumor progression, the development of metastases, and poor survival.^{20,21} Whether or not DNA ploidy is an independent variable remains questionable. Assessing the impact of DNA ploidy changes in early-stage disease is complicated by the long survival times of these patients. In patients with pT2 disease,

In addition to the amount of DNA present in the nucleus, measurements of nuclear size and shape have also been reported to be of prognostic significance.

Gleason score and pre-operative PSA remain the most effective tool for predicting prognosis. However, with respect to pT3 tumors, DNA ploidy has prognostic value in predicting response to hormonal therapy.²² Results on metastatic prostate cancer remain controversial.^{23,24} Although recent attempts have been made to evaluate DNA ploidy in material from needle biopsies, it is likely that sampling errors may confound these attempts and preclude clinical utility.

In addition to the amount of DNA present in the nucleus, measurements of nuclear size and shape (termed

approaches. These include mitotic figure counting, static and flow cytometry, bromodeoxyuridine labeling, and immunostaining with antibodies directed against components of proliferating cells, such as proliferating cell nuclear antigen (PCNA), Ki-67, p21, and cyclin D1.²⁹⁻³¹ Essentially all of these studies have revealed very low rates of proliferation in most tumors. Attempts have also been made to correlate proliferative activity with disease outcome.^{32,33} Although most studies have indicated strong correlation between features such as proliferation and histologic

grade and/or tumor stage, there are only very limited data to suggest an independent prognostic significance associated with measurements of proliferation indices.

One of the hallmark features of lethal malignancies is their ability to invade and/or metastasize. Obligate

$\alpha 5$, αv and $\beta 4$ have been reported. Higher expression of $\alpha 6$ is correlated with increased invasion.³⁸ The $\beta 1 C$ variant, reported to inhibit tumor growth, is down-regulated in prostate cancer.³⁹ E-cadherin is a well-known member of another large family of proteins associated with calcium-

reported to occur in the majority of prostate cancers.⁴³ Decreased expression was also found to correlate with high-grade and DNA aneuploidy.⁴⁴ To the contrary, other authors have reported that increased CD44 expression correlates with high grade and advanced stage. Clearly, additional studies are necessary to resolve these differences.

There are only very limited data to suggest an independent prognostic significance associated with measurements of proliferation indices.

to this process is the loss of intercellular adhesion that normally holds epithelial cells together at their sites of origin. Within normal tissues, epithelial cells have a variety of mechanisms that attach them to neighboring cells and to their basement membrane. Assessing the status of mechanisms by which cells adhere is probably important for both providing biologic insight as well as developing prognostic markers. Some studies have suggested that an inverse relation exists between the down-regulation of selected cell adhesion molecules in high-grade prostatic intraepithelial neoplasia (HGPIN) and prostate cancer and the up-regulation of invasion-related enzymes believed to be involved in the degradation of basement membrane, such as collagenases and matrix metalloproteinases.^{34,35} The adhesion molecules that have attracted the most attention are the integrins, the cadherins, C-CAM, and CD44. The integrins are a large family of heterodimeric plasma membrane receptors composed of alpha and beta subunits. The patterns and combinations of subunits that are expressed appear to be different in normal epithelium, HGPIN, and carcinoma. In general, loss of expression is associated with the malignant phenotype and tumor progression.^{36,37} Specifically, losses of integrins $\alpha 2$, $\alpha 4$,

dependent adhesion. Several investigators have indicated that a correlation exists between reduction or loss of E-cadherin expression and advanced-stage, higher-grade prostate cancer. Although one third of organ-confined prostate cancers exhibit abnormal E-cadherin expression, more than 75% of metastatic carcinomas have this abnormality.⁴⁰ C-CAM is a calcium-independent molecule that actually belongs to the immunoglobulin family of genes. Its sequence is also similar to carcinoem-

Vitamin D and Prostate Cancer: From Laboratory to Bedside

Although it is recognized that prostatic cancer is an androgen-dependent disease, it is now very apparent that a variety of other steroid and peptide hormones are capable of regulating the growth and differentiation of prostatic cancer cells. Early epidemiologic studies have indicated that the clinical emergence of prostate cancer may be at least partially explained by vitamin D deficiency.⁴⁵ In addition to age and race, these studies linked geographic location to prostate cancer mortality through potential actions on vitamin D levels. The link

Several investigators have indicated that a correlation exists between reduction or loss of E-cadherin expression and advanced-stage, higher-grade prostate cancer.

bryonic antigen. Decreased expression has been reported in HGPIN, and its expression was not detected in carcinomas.⁴¹ This suggests that C-CAM is a possible suppressor gene that is altered early in prostatic oncogenesis. CD44 is another transmembrane protein believed to be involved with cell-cell and cell-matrix interactions. A soluble form of CD44 known as variant 5 (v5) was found to be lower in the serum of patients with prostate cancer or benign prostatic hyperplasia (BPH) than controls.⁴² However, down-regulation of the CD44 standard isoform has been

between latitude and vitamin D action was clear, because ultraviolet light is required for the initial conversion of the vitamin D precursor 7-dehydrocholesterol to vitamin D₃. What remained to be shown was that there was a biologic basis for the remainder of this hypothesis.

Vitamin D is known to act through both nuclear receptors (the genomic pathway) and cell membrane receptors (the non-genomic pathway). Vitamin D receptors had been found in various non-prostate benign and malignant epithelial cells. Using nuclear receptor binding assays,

Miller et al were the first to show that specific, high-affinity binding sites for the active metabolite of vitamin D ($1\alpha,25$ -dihydroxyvitamin D₃) are present in prostate cancer cells.⁴⁶ The number of nuclear vitamin D receptors per cell varies widely among different cell lines ranging from approximately 500 to 18,000.^{47,48}

It is known that vitamin D can have a marked antiproliferative effect of various non-prostate malignant cell lines in vitro.

Recently, it has been found that many of the prostate cancer cell lines that were used for such studies actually represent contaminants from PC-3. This finding has led to the interesting observation that ALVA-31 cells, which is now known to be a subline of PC-3, have much higher expression of vitamin D receptors than the parental line. The mechanisms through which this increased expression has occurred are not presently known. If such a change could be induced, however, this could prove to be a valuable therapeutic approach.

It is known that vitamin D can have a marked antiproliferative effect of various non-prostate malignant cell lines in vitro. This has also been found in various prostatic carcinoma cell lines. However, it is interesting to note that the degree of antiproliferative effect varies widely between cell lines. This suggests that whereas some cells remain sensitive to the effects of vitamin D, others may have become resistant. The mechanisms through which cells become vitamin D resistant are also currently under study and may provide critical insights into the use of vitamin D in the therapy of prostate cancer.

Treatment of patients with high doses of vitamin D would likely be stymied by the hypercalcemic effects

of this compound. To date, however, hypercalcemic effects in treated prostate cancer patients have been relatively inconsequential.⁴⁵ In addition, numerous synthetic vitamin D analogues have been developed that have increased pro-differentiative effects while displaying a lower hypercalcemic potential. Studies at

the University of Colorado in prostate cancer cells have indicated that those analogues with hexafluorinated side chains and increased saturation of the bonds in this chain have a greater antiproliferative effect on prostate cancer cells than does the parent molecule.⁴⁹ Clinical trials with these compounds are attractive but are currently not possible owing to lack of appropriate agents approved for use in humans.

Studies with vitamin D and its analogues have also made it clear that effects on differentiation accompany the antiproliferative effects. Using the well-characterized prostate cancer cell line LNCaP, University of Colorado researchers have found that vitamin D and its analogues increase the synthesis and release of both PSA and prostate-specific acid phosphatase into the culture medium. This finding is of great clinical potential, because it indicates that in clinical trials, an early increase in a patient's serum PSA levels might indicate success rather than failure.

The mechanisms through which vitamin D induces a decrease in proliferation are now becoming clear. Preliminary results indicate that an increase in the cell cycle mediator p21 appears to be necessary for the antiproliferative effects. This

increase occurs within days and could be caused by indirect actions on other genes. Consistent with this possibility, it has also been found that in some cells, but not others, p21 induction by vitamin D appears to require a prior induction of transforming growth factor beta. Again, the mechanisms of this induction, and alternate pathways will probably offer clues to methods for the appropriate use of these compounds in the management of clinical disease.

Finally, it has been found in studies on prediagnostic sera levels that decreased levels of vitamin D metabolites accompany both the presence of prostate cancer and its biologic aggression. More interestingly, it is now known that as many as 37% of hospitalized patients whose vitamin D intake is greater than the current recommended daily allowance can be vitamin D deficient. It also appears that vitamin D deficiency may be particularly common among men with hormone-refractory or advanced-stage prostate cancer. It would seem reasonable, therefore, to include vitamin D supplementation into treatment strategies for these patients. To this end, the ability of vitamin D to potentiate the effects of chemotherapeutic agents, not otherwise found to be of great value in the treatment of prostate cancer, has been examined. Vitamin D can have a synergistic effect on growth arrest of prostate cancer cells induced by cisplatin or carboplatin.⁵⁰ These findings have led to a clinical trial, in which patients are supplemented with vitamin D prior to receiving carboplatin therapy. Patients are also given dexamethasone prior to either of these agents. The results are preliminary but indicate that as many as 70% of men will respond to such therapy with decreases in their PSA of more than 50% over prolonged periods of time.

These findings indicate the potential for vitamin D and perhaps other forms of differentiation therapy to contribute to our armamentarium for the treatment of prostate cancer.

The Insulin-Like Growth Factor System in the Control of Carcinoma of the Prostate

The insulin-like growth factor (IGF) system is a complex group of molecules composed of three ligands (insulin, IGF-I, and IGF-II) and seven known binding proteins that can either enhance or inhibit the effects of the ligands.⁵¹ The system has been studied in carcinomas of the breast, sarcomas, Wilms' tumors, and colorectal cancers.⁵² Early studies on the role of this system in prostate cancer demonstrated that levels of binding protein 2 were increased at the same time that levels of binding protein 3 were decreased in the serum of patients with advanced-stage disease.⁵³ At the tissue level, it was next shown that the expression of binding protein 2 mRNA and protein were increased in carcinomas.⁵⁴ However, although the levels of binding proteins 3 and 4 were decreased, the levels of their mRNA expression had remained constant, suggesting the regulation of their protein levels was post-translational.⁵⁴ This observation is important, because protein 2 contains an RGD sequence and can increase cell motility, whereas binding proteins 3 and 4 are growth inhibitory.⁵⁴ An increase in protein 2 accompanied by decreases in proteins 3 and 4 could cause cells to proliferate and become more metastatic.⁵⁴

The next advance in these studies came through the development of a series of cell lines derived from benign prostatic epithelial cells infected with SV40 large T antigen. A series of cell lines, including p-69, M-2182, and M-12 were established by passage as xenografts.⁵⁵ Although

p-69 is poorly tumorigenic and non-metastatic, the M-12 line is 100% tumorigenic and metastatic when implanted either orthotopically or intraperitoneally. Transfections were used to create an IGF type-I receptor over-expressing cell, called M-12 LISN, and its receptor-deficient counterpart, M-12 LNL-6 cells. Radioligand binding and Scatchard analysis revealed that the LISN cells contained significantly more recep-

tors per cell than the LNL-6 cells. Soft agar assays for anchorage-independent growth revealed that LISN cells formed 75% fewer colonies than LNL-6 cells, suggesting that expression of receptors should decrease the tumorigenicity of these cells. Consistent with this postulate, the volumes of tumors formed in xenografts with LISN cells were 6-fold less than those derived from LNL-6 cells. p-69 cells were also found to contain very small amounts of telomerase activity, whereas M-12 cells contained approximately 100-fold higher levels. However, when the transfectants were examined, LISN cells were found to contain 10-fold less telomerase than LNL-6 cells. This suggests that the LISN cells are more terminally differentiated but at the same time more responsive to growth factors. The results of these studies could form the basis for a specific approach to gene therapy, in which either the expression of the IGF receptors or the expression of IGFs themselves are modified.

With respect to human applications of this research, six attempts have been made to correlate serum IGF-I levels with the risk for developing prostate cancer.⁵⁶ Three of these have

shown no association, and three have shown association. The interesting aspect of this conundrum is that different assays were used to obtain the results between the different trials. It would seem that until there is better standardization of assays and larger numbers of patients can be examined, we will not be able to accurately assess the true relations of this pathway. However, studies by Wolk et al⁵⁷ on IGFBP-3 and IGF-I levels in the

As many as 37% of hospitalized patients whose vitamin D intake is greater than the current recommended daily allowance can be vitamin D deficient.

serum of prostate cancer patients may provide some additional insight. These workers found that in patients who had serum PSA levels of 3 or less, the presence of an elevated level of IGF-I and lower levels of IGFBP-3 were indicative of the presence of cancer. Thus, although it is not yet clear that the IGF axis is useful to diagnose cancer by itself, it may be of assistance in clarifying otherwise ambiguous serum PSA results.

What's New in Radiologic Staging?

Much of our approach to clinical management of prostate cancer patients is highly dependent on accurate clinical staging. Although a number of surrogate markers, such as serum PSA levels and Gleason scores, have helped in assessing and predicting stage, they are not always as accurate as we would desire. Ultimately, the preferred approach to staging would be an actual visualization of the whereabouts of locally invasive or metastatic prostate cancer, such that the stage of patients could be objectively defined rather than predicted.^{58,59}

At present, radionuclide bone scans remain the most effective way

to screen the entire skeleton for metastases in high-risk patients. Once abnormalities are detected, additional evaluation may be necessary using computed tomography (CT)

clinical use and a relatively large experience with it has accumulated.⁶⁴⁻⁶⁹ Because of limited experience in controlled trials, however, its precise role in clinical management is

In patients with a high clinical suspicion of extraprostatic disease and no evidence of distant metastases, MRI remains a reasonable approach to evaluate for adenopathy or local extension of carcinoma.

or magnetic resonance imaging (MRI). In addition, conventional x-rays can often be used to confirm the nature of suspicious areas from bone scans. CT and MRI can also be used to evaluate patients for abdominal or pelvic lymphadenopathy. The relatively limited sensitivity of these studies has led to evaluation of MRI spectroscopy, indium-111 capromab pentetide (ProstaScint), and positron emission tomography (PET) for this purpose. At present, however, the use of these alternate approaches remains experimental.

The use of color and/or power Doppler ultrasound may increase the sensitivity of detection of primary lesions in the prostate, but reported results have been variable.⁶⁰⁻⁶² Ultrasound contrast agents are currently under investigation. Initial studies with these agents suggest that contrast agents improve sonographic accuracy both in detection and in determining the extent of local invasion.⁶³ Other technical advances including harmonic imaging and three-dimensional ultrasound are also being evaluated.

The use of radiolabeled antibodies directed against proteins on the surfaces of malignant cells has long been investigated as a method to detect small quantities of those cells anywhere in the body. Several different antibodies have been used to evaluate prostatic carcinomas. At present, ProstaScint is approved for

not yet well defined. Situations in which some practitioners find it of clinical utility include: 1) evaluation of residual or recurrent disease after local treatment including prostatectomy; and 2) providing prognostic information or facilitating treatment planning at the time of initial diagnosis. However, many find the limited spatial resolution in these images as well as the high background associated with this agent to be unacceptable.

MRI, especially using an endorectal coil, is the most accurate cross-sectional imaging method available to evaluate the local extent of prostate carcinoma invasion.⁷⁰⁻⁷⁶ Because of its cost and limitations, however, it cannot be universally recommended. In patients with a high clinical suspicion of extraprostatic disease and no evidence of distant metastases, MRI remains a reasonable approach to evaluate for adenopathy or local extension of carcinoma.

PET imaging offers the potential to detect malignant cells based on their increased glucose metabolism wherever they are located within the body.⁷⁷⁻⁸² The potential exists to use PET in the pretreatment search for metastases to the bone or lymph nodes. This technique could also be applied to the detection of recurrences after various forms of local treatment. PET appears to be very useful for determining the extent of disease with several nonprostatic malignancies. However, some pre-

liminary studies in patients with prostate cancer have been disappointing. Its use at present should, therefore, remain investigational.

One additional imaging modality should also be considered. Proton magnetic resonance spectroscopy (MR spectroscopy) can be used to evaluate tissue metabolic activity as reflected by local concentrations of metabolic breakdown products.⁸³⁻⁸⁵ One group of investigators has found that spectroscopy added to conventional rectal coil MRI improves the accuracy in determining both the intraprostatic location and extent of involvement by carcinoma.⁸⁶ Evidence has also been presented indicating that MR spectroscopy can improve accuracy in the diagnosis of extracapsular disease. At present, however, the technique remains largely experimental and is not widely available. ■

References

1. Christensson A, Laurell CB, Lilja H. Enzymatic activity of prostate-specific antigen and its reaction with extracellular serine proteinase inhibitors. *Eur J Biochem.* 1990;194:755-763.
2. Stenman U, Leinonen J, Alfthan H, et al. A complex between PSA and alpha-1-antichymotrypsin is the major form of PSA in serum of patients with prostatic cancer: assay of the complex improves clinical sensitivity for cancer. *Cancer Res.* 1991;51:222-226.
3. Espana F, Sanchez CJ, Vera CD, et al. A quantitative ELISA for the measurement of complexes of prostate-specific antigen with protein C inhibitor when using a purified standard. *J Lab Clin Med.* 1993;122:711-719.
4. Christensson A, Lilja H. Complex formation between protein C inhibitor and prostate-specific antigen in vitro and in human semen. *Eur J Biochem.* 1994;220:45-53.
5. Christensson A, Bjork T, Nilsson O, et al. Serum prostate-specific antigen complexed to alpha 1-antichymotrypsin as an indicator of prostate cancer. *J Urol.* 1993;150:100-105.
6. Luderer AA, Chen Y, Thiel R, et al. Measurement of the proportion of free to total PSA improves diagnostic performance of PSA in the diagnostic gray zone of total PSA. *Urology.* 1995;46:187-194.
7. Higashihara E, Nutahara K, Kojima M, et al. Significance of serum free prostate-specific antigen in the screening of prostate cancer. *J Urol.* 1996;156:1964-1968.
8. Catalona WJ, Partin AW, Slawin KM, et al. Use of the percentage of free prostate-specific antigen to enhance differentiation of prostate cancer from benign prostatic disease: a prospective multicenter clinical trial. *JAMA.* 1998;279:1542-1547.
9. Piironen T, Pettersson K, Suonpaa M, et al. In vitro stability of free prostate-specific antigen

- (PSA) and prostate-specific antigen (PSA) complexed to alpha-1-antichymotrypsin in blood samples. *Urology*. 1996;48:81-87.
10. Woodrum DL, Brawer MK, Partin AW, et al. Interpretation of free prostate-specific antigen: clinical research studies for the detection of prostate cancer. *J Urol*. 1998;159:5-12.
 11. Roth HJ, Christensen-Stewart S, Brawer MK. A comparison of three free and total PSA assays. *PCPD*. 1998;1:326-331.
 12. Nixon RG, Gold MH, Blase AB, et al. Comparison of three investigative assays for the free form of prostate-specific antigen. *J Urol*. 1998;160:420-425.
 13. Allard WJ, Zhou Z, Yeung KK. Novel immunoassay for the measurement of complexed prostate-specific antigen in serum. *Clin Chem*. 1998;44:1216-1223.
 14. Brawer MK, Meyer GE, Letran JL, et al. Measurement of complexed PSA improves specificity for early detection of prostate cancer. *Urology*. 1998;52:372-378.
 15. Brawer MK, Cheli CD, Neaman IE, et al. Complexed prostate specific antigen provides significant enhancement of specificity compared with total prostate specific antigen for detecting prostate cancer. *J Urol*. 2000;163:1476-1480.
 16. Simon R, Altman DGT. Statistical aspects of prognostic factor studies in oncology. *Br J Cancer*. 1994;69:979-985.
 17. Hunter DJ. Methodological issues in the use of biological markers in cancer epidemiology: cohort studies. *LARC Sci Publ*. 1997;142:39-46.
 18. Shankey TV, Kallioniemi OP, Koslowski JM, et al. Consensus review of the clinical utility of DNA content cytometry in prostate cancer. *Cytometry*. 1993;14:497-500.
 19. Centeno BA, Zietman AL, Shipley WU, et al. Flow cytometric analysis of DNA ploidy, percent S-phase fraction, and total proliferative fraction as prognostic indicators of local control and survival following radiation therapy for prostate carcinoma. *Int J Radiation Oncology Biol Phys*. 1994;30:309-315.
 20. Carmichael MJ, Veltri RW, Partin AW, et al. Deoxyribonucleic acid ploidy analysis as a predictor of recurrence following radical prostatectomy for stage T2 disease. *J Urol*. 1995;153:1015-1019.
 21. Hawkins CA, Bergstralh EJ, Lieber MM, et al. Influence of DNA ploidy and adjuvant treatment on progression and survival in patients with pathologic stage T3 (PT3) prostate cancer after radical retropubic prostatectomy. *Urology*. 1995;46:356-364.
 22. Hayashi T, Mouri J, Kurosu S, et al. Flow cytometric analysis of prostatic adenocarcinomas after hormone therapy. *Nippon Hinyokika Gakkai Zasshi*. 1994;85:339-245.
 23. Jorgensen T, Kanagasigam Y, Kaalhus O, et al. Prognostic factors in patients with metastatic (stage D2) prostate cancer: experience from the Scandinavian Prostate Cancer Group Study-2. *J Urol*. 1997;158:164-170.
 24. Peters-Gee JM, Miles BJ, Cerny JC, et al. Prognostic significance of DNA quantitation in stage D1 prostate carcinoma with the use of image analysis. *Cancer*. 1992;70:1159-1165.
 25. Vesalainen S, Lipponen P, Talja M, et al. Nuclear morphometry is of independent prognostic value only in T1 prostatic adenocarcinomas. *Prostate*. 1995;27:110-117.
 26. Kanamaru H, Zhang YH, Takahashi, et al. Analysis of the mechanism of discrepant nuclear morphometric results comparing pre-operative biopsy and prostatectomy specimens. *Urology*. 2000;56:342-345.
 27. Zhang YH, Kanamaru H, Oyama N, et al. Prognostic value of nuclear morphometry on needle biopsy from patients with prostate cancer: is volume-weighted mean nuclear volume superior to other morphometric parameters? *Urology*. 2000;55:377-381.
 28. Bacus JW, Bacus JV, Stoner GD, et al. Quantitation of preinvasive neoplastic progression in animal models of chemical carcinogenesis. *J Cell Biochem Suppl*. 1997;28-29:21-38.
 29. Drobniak M, Osman I, Scher HI, et al. Overexpression of cyclin D1 is associated with metastatic prostate cancer to bone. *Clin Cancer Res*. 2000;6:1891-1895.
 30. Kaibuchi T, Furuya Y, Akakura K, et al. Changes in cell proliferation and apoptosis during local progression of prostate cancer. *Anticancer Res*. 2000;20:1135-1139.
 31. Mucci NR, Rubin MA. Expression of nuclear antigen Ki-67 in prostate cancer needle biopsy and radical prostatectomy. *J Natl Cancer Inst*. 2000;92:1941-1942.
 32. Cheng L, Pisanksky TM, Sebo TJ, et al. Cell proliferation in prostate cancer patients with lymph node metastasis: a marker for progression. *Clin Cancer Res*. 1999;5:2820-2823.
 33. Keshgegian AA, Johnston E, Canaan A, et al. Bcl-2 oncoprotein positivity and high MIB-1 (ki-67) proliferative rate are independent predictive markers for recurrence in prostate carcinoma. *Am J Clin Pathol*. 1998;110:443-449.
 34. Upadhyay J, Shekarriz B, Nemeth JA, et al. Membrane type 1-matrix metalloproteinase (MT1-MMP) and MMP-2 immunolocalization in human prostate: change in cellular localization associated with high-grade prostatic intraepithelial neoplasia. *Clin Cancer Res*. 1999;5:4105-4110.
 35. Montironi R, Mazzucchelli R, Stramazzotti D, et al. Expression of pi-class glutathione S-transferase: two populations of high grade prostatic intraepithelial neoplasia with different relations to carcinoma. *Mol Pathol*. 2000;53:122-128.
 36. Allen MV, Smith GJ, Juliano R, et al. Downregulation of the beta4 integrin subunit in prostatic carcinoma and prostatic intraepithelial neoplasia. *Hum Pathol*. 1998;29:311-318.
 37. Zheng DG, Woodward AS, Fornaro M, et al. Prostatic carcinoma cell migration via alpha(v)beta3 integrin is modulated by a focal adhesion kinase pathway. *Cancer Res*. 1999;59:1655-1664.
 38. Cress AE, Rabinovitz I, Zhu W, et al. The alpha 6 beta 1 and alpha 6 beta 4 integrins in human prostate cancer progression. *Cancer Metastasis Rev*. 1995;14:219-228.
 39. Fornaro M, Manzotti M, Tallini G, et al. Beta 1C

Main Points

- Much effort has been expended in attempting to develop specific assays for the complexed form of PSA; one company has developed a specific assay for complex PSA that has been shown to be highly reproducible as well as specific.
- Because the complexed PSA assay is equivalent to total PSA for monitoring or staging disease, it would thus appear to be the best possible form of PSA assay for all clinical uses.
- Many studies have indicated that abnormal DNA content in tumor cells (DNA ploidy) correlates with tumor progression, the development of metastases, and poor survival; whether or not DNA ploidy is an independent variable remains questionable.
- Some studies have suggested that an inverse relation exists between the down-regulation of selected cell adhesion molecules (eg, the integrins, the cadherins, C-CAM, and CD44) in prostate cancer and the up-regulation of invasion-related enzymes believed to be involved in the degradation of basement membrane.
- Early epidemiologic studies have indicated that the clinical emergence of prostate cancer may be at least partially explained by vitamin D deficiency.
- Although it is not yet clear that the insulin-like growth factor axis is useful to diagnose cancer by itself, it may be of assistance in clarifying otherwise ambiguous serum PSA results.
- At present, radionuclide bone scans remain the most effective way to screen the entire skeleton for metastases in high-risk patients; once abnormalities are detected, additional evaluation may be necessary using CT or MRI.
- PET imaging offers the potential to detect malignant cells based on their increased glucose metabolism wherever they are located within the body; however, some preliminary studies in patients with prostate cancer have been disappointing. Use of PET at present should, therefore, remain investigational.

- integrin in epithelial cells correlates with a nonproliferative phenotype: forced expression of beta1C inhibits prostate epithelial cell proliferation. *Am J Pathol*. 1998;153:1079-1087.
40. Umbas R, Issacs WB, Brinquier PP, et al. Decreased E-Cadherin expression is associated with poor prognosis in patients with prostate cancer. *Cancer Res*. 1994;54:3929-3933.
 41. Kleinerman DI, Troncoso P, Lin SH, et al. Consistent expression of an epithelial cell adhesion molecule (C-CAM) during human prostate development and loss of expression in prostate cancer: implication as a tumor suppressor. *Cancer Res*. 1995;55:1215-1220.
 42. Aaltomaa S, Lipponen P, Viitanen J, et al. Prognostic value of CD44 standard, variant isoforms 3 and 6 and catenin expression in local prostate cancer treated by radical prostatectomy. *Eur Urol*. 2000;38:555-562.
 43. Kallakury BV, Sheehan CE, Ambros RA, et al. Correlation of p34cdc2 cyclin-dependent kinase overexpression, CD44s downregulation, and HER-2/neu oncogene amplification with recurrence in prostatic adenocarcinomas. *J Clin Oncol*. 1998;16:1302-1309.
 44. Noordzij MA, van Steenbrugge GJ, Schroder FH, et al. Decreased expression of CD44 in metastatic prostate cancer. *Int J Cancer*. 1999;84:478-483.
 45. Miller GJ. Vitamin D and prostate cancer: biologic interactions and clinical potentials. *Cancer Metastasis Rev*. 1999;17:353-360.
 46. Miller GJ, Stapleton GE, Ferrara JA, et al. The human prostate carcinoma cell line LNCaP expresses biologically active, specific receptors for 1a,25 dihydroxyvitamin D3. *Cancer Res*. 1992;52:515-520.
 47. Miller GJ, Stapleton GE, Hedlund TE, Moffatt KA. Vitamin D receptor expression, 24-hydroxylase activity, and inhibition of growth by 1a,25-dihydroxyvitamin D3 in seven human prostatic carcinoma cell lines. *Clin Cancer Res*. 1995;1:997-1003.
 48. Hedlund TE, Moffatt KA, Miller GJ. Stable overexpression of the nuclear vitamin D receptor in the human prostatic carcinoma cell line JCA-1: evidence that the antiproliferative effects of 1a,25-dihydroxyvitamin D3 are mediated exclusively through the genomic signaling pathway. *Endocrinology*. 1996;137:1554-1561.
 49. Hedlund TE, Moffatt KA, Uskokovic MR, Miller GJ. Three synthetic vitamin D analogues induce prostate-specific acid phosphatase and prostate-specific-antigen, while inhibiting the growth of human prostate cancer cells in a vitamin D receptor-dependent fashion. *Clin Cancer Res*. 1997;3:1331-1338.
 50. Moffatt KA, Johannes WU, Miller GJ. 1a,25 dihydroxyvitamin D3 and platinum drugs act synergistically to inhibit the growth of prostate cancer cell lines. *Clin Cancer Res*. 1999;5:695-703.
 51. Grimberg A, Rajah R, Zhao H, Cohen P. The prostatic IGF system: new levels of complexity. In: Takano K, Hizuka N, Takahashi SI, eds. *Molecular Mechanisms to Regulate the Activities of Insulin-Like Growth Factors*. Amsterdam: Elsevier; 1998:205.
 52. LeRoith D, Baserga R, Helman L, Roberts CT. Insulin-like growth factors and cancer. *Ann Intern Med*. 1995;122:54-59.
 53. Kanety H, Madjar Y, Dagan Y, et al. Serum insulin-like growth factor-binding protein-2 (IGFBP-2) is increased and IGFBP-3 is decreased in patients with prostate cancer: correlation with serum prostate-specific antigen. *J Clin Endocrinol Metab*. 1993;77:229-233.
 54. Thrasher JB, Tennant MK, Twomey PA, et al. Immunohistochemical localization of insulin-like growth factor binding proteins 2 and 3 in prostate tissue: clinical correlations. *J Urol*. 1996;115:999-1003.
 55. Plymate SR, Tennant M, Birmbaum RS et al. The effect of the insulin-like growth factor system in human prostate epithelial cells of immortalization and transformation by Simian Virus-40 T antigen. *J Clin Endocrinol Metab*. 1996;81:3709-3716.
 56. Grimberg A, Cohen P. Growth hormone and prostate cancer: guilty by association. *J Endocrinol Invest*. 1999;22:64-73.
 57. Wolk A, Anderson SO, Mantzoros CS, et al. Can measurements of IGF-1 and IGFBP-3 improve the sensitivity of prostate-cancer screening? *Lancet*. 2000;356:1902-1904.
 58. Yu KK, Hawkins RA. The prostate: diagnostic evaluation of metastatic disease. *Radiol Clin N Am*. 2000;38:139-157.
 59. Yu KK, Hricak H. Imaging prostate cancer. *Radiol Clin N Am*. 2000;38:59-85.
 60. Cho JY, Kim SH, Lee SE. Diffuse prostatic lesions: role of color Doppler and power Doppler ultrasonography. *J Ultrasound Med*. 1998;17:283-287.
 61. Cornud F, Belin X, Piron D, et al. Color doppler-guided prostate biopsies in 591 patients with an elevated serum PSA level: impact on Gleason score for nonpalpable lesions. *Urology*. 1997;49:709-715.
 62. Ismail M, Petersen RO, Alexander AA, et al. Color doppler imaging in predicting the biologic behavior of prostate cancer: correlation with disease-free survival. *Urology*. 1997;50:906-912.
 63. Halpern EJ, Verkh L, Forsberg F, et al. Initial experience with contrast-enhanced sonography of the prostate. *AJR Am J Roentgenol*. 2000;174:1575-1580.
 64. Gregorakis AK, Holmes EG, Murphy GP. Prostate-specific membrane antigen: current and future utility. *Semin Urol Oncol*. 1998;16:2-12.
 65. Kahn D, Williams RD, Manyak MJ, et al. 111-Indium-capromab pendetide in the evaluation of patients with residual or recurrent prostate cancer after radical prostatectomy. *J Urol*. 1998;159:2041-2047.
 66. Levesque PE, Nieh PT, Zinman LN, et al. Radiolabeled monoclonal antibody indium 111-labeled CYT-356 localizes extraprostatic recurrent carcinoma after prostatectomy. *Urology*. 1998;51:978-984.
 67. Manyak MJ, Javitt MC. The role of computerized tomography, magnetic resonance imaging, bone scan, and monoclonal antibody nuclear scan for prognosis prediction in prostate cancer. *Semin Urol Oncol*. 1998;16:145-152.
 68. Murphy GP, Maguire RT, Rogers B, et al. Comparison of serum PSMA, PSA levels with results of Cytogen-356 ProstaScint scanning in prostatic cancer patients. *Prostate*. 1997;33:281-285.
 69. Quintana JC, Blend MJ. The dual-isotope ProstaScint imaging procedure: clinical experience and staging results in 145 patients. *Clin Nucl Med*. 2000;25:33-40.
 70. D'Amico AV, Schnall M, Whittington R, et al. Endorectal coil magnetic resonance imaging identifies locally advanced prostate cancer in select patients with clinically localized disease. *Urology*. 1998;51:449-454.
 71. D'Amico AV. Re: endo-rectal coil magnetic resonance imaging in clinically localized prostate cancer: is it accurate? *J Urol*. 1997;157:1371.
 72. Getty DJ, Seltzer SE, Tempany CM, et al. Prostate cancer: relative effects of demographic, clinical, histologic, and MR imaging variable on the accuracy of staging. *Radiology*. 1997;204:471-479.
 73. Ikonen S, Karkkainen P, Kivisaari L, et al. Magnetic resonance imaging of clinically localized prostatic cancer. *J Urol*. 1998;159:915-919.
 74. Silverman JM, Krebs TL. MR imaging evaluation with a transrectal surface coil of local recurrence of prostatic cancer in men who have undergone radical prostatectomy. *AJR Am J Roentgenol*. 1997;168:379-385.
 75. Tempany CM, Langlotz CP. Re: endo-rectal coil magnetic resonance imaging in clinically localized prostate cancer: is it accurate? *J Urol*. 1997;157:1371-1372.
 76. Yu KK, Hricak H, Alagappan R, et al. Detection of extracapsular extension of prostate carcinoma with endorectal and phased-array coil MR imaging: multivariate feature analysis. *Radiology*. 1997;202:697-702.
 77. Effert PJ, Handts RB, Wolff JM, Bull U. Metabolic imaging of untreated prostate cancer by positron emission tomography with [18]fluorine-labeled deoxyglucose. *J Urol*. 1996;155:994-998.
 78. Haseman NK, Reed NL, Rosenthal SA. Monoclonal antibody imaging of occult prostate cancer in patients with elevated prostate-specific antigen: positron emission tomography and biopsy correlation. *Clin Nucl Med*. 1996;21:703-713.
 79. Heicappell R, Muller-Mattheis V, Reinhardt M, et al. Staging of pelvic lymph nodes in neoplasms of the bladder and prostate by positron emission tomograph with 2-[(18)F]-2-deoxy-D-glucose. *Eur Urol*. 1999;36:582-587.
 80. Hofer C, Laubenbacher C, Block T, et al. Fluorine-18-fluorodeoxyglucose positron emission tomography is useless for the detection of local recurrence after radical prostatectomy. *Eur Urol*. 1999;36:31-35.
 81. Schirmer H, Guhlmann A, Elsner K, et al. Sensitivity in detecting osseous lesions depends on anatomic localization: planar bone scintigraphy versus 18F PET. *J Nucl Med*. 1999;40:1623-1629.
 82. Seltzer MA, Barbaric Z, Beldegrun A, et al. Comparison of helical computerized tomography, position emission tomography and monoclonal antibody scans for evaluation of lymph node metastases in patients with prostate specific antigen relapse after treatment for localized prostate cancer. *J Urol*. 1999;162:1322-1328.
 83. Kurhanewicz J, Vigneron DB, Males RG, et al. The prostate: MR imaging and spectroscopy. Present and future. *Radiol Clin N Am*. 2000;38:115-138.
 84. Scheidler J, Hricak H, Vigneron DB, et al. Prostate cancer: localization with three-dimensional proton MR spectroscopic imaging-clinical-pathologic study. *Radiology*. 1999;213:473-480.
 85. Wefer AE, Hricak H, Vigneron DB, et al. Sextant localization of prostate cancer: comparison of sextant biopsy, magnetic resonance imaging and magnetic resonance spectroscopic imaging with step section histology. *J Urol*. 2000;164:400-404.
 86. Yu KK, Scheidler J, Hricak H, et al. Prostate cancer: prediction of extracapsular extension with endorectal MR imaging and three-dimensional proton MR spectroscopic imaging. *Radiology*. 1999;213:481-488.